

EXHIBIT E

EXHIBIT E

MEDLINE(R)

(c) format only 2006 Dialog. All rights reserved.

08202489 PMID: 2677748

Binding activities of a repertoire of single immunoglobulin variable domains secreted from *Escherichia coli*.

Ward E S; Gussow D; Griffiths A D; Jones P T; Winter G

MRC Laboratory of Molecular Biology, Cambridge, UK.

Nature (ENGLAND) Oct 12 1989 , 341 (6242) p544-6 , ISSN: 0028-0836--

Print Journal Code: 0410462

Publishing Model Print; Comment in Nature. 1989 Oct 12;341(6242) 484-5;

Comment in PMID 2677746

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: MEDLINE; Completed

In antibodies, a heavy and a light chain variable domain, VH and VL, respectively, pack together and the hypervariable loops on each domain contribute to binding antigen. We find, however, that isolated VH domains with good antigen-binding affinities can also be prepared. Using the polymerase chain reaction, diverse libraries of VH genes were cloned from the spleen genomic DNA of mice immunized with either lysozyme or keyhole-limpet haemocyanin. From these libraries, VH domains were expressed and secreted from *Escherichia coli*. Binding activities were detected against both antigens, and two VH domains were characterized with affinities for lysozyme in the 20 nM range. Isolated variable domains may offer an alternative to monoclonal antibodies and serve as the key to building high-affinity human antibodies. We suggest the name 'single domain antibodies (dAbs)' for these antigen binding demands.